



Genome-Enabled Molecular Tools for Reductive Dehalogenation

- A Shift in Paradigm for Bioremediation -

Alfred M. Spormann
Departments of Chemical and of Civil & Environmental Engineering
Stanford University

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to maintaining the data needed, and completing and reviewing the collect including suggestions for reducing this burden, to Washington Headqu VA 22202-4302. Respondents should be aware that notwithstanding at does not display a currently valid OMB control number.	tion of information. Send comments relaters Services, Directorate for Information	regarding this burden estimate of mation Operations and Reports	or any other aspect of the s, 1215 Jefferson Davis	nis collection of information, Highway, Suite 1204, Arlington	
1. REPORT DATE NOV 2011	2. REPORT TYPE		3. DATES COVE 00-00-2011	RED 1 to 00-00-2011	
4. TITLE AND SUBTITLE			5a. CONTRACT	NUMBER	
Genome-Enabled Molecular Tools for Reductive Dehalogenation		nation	5b. GRANT NUMBER		
		5c. PROGRAM ELEMENT NUMBER			
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Stanford University, Departments of Chemical Engineering, and of Civil & Environmental Engineering, 318 Campus Drive, Stanford, CA, 94305			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/M NUMBER(S)	ONITOR'S REPORT	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution	ion unlimited				
13. SUPPLEMENTARY NOTES Presented at the Partners in Environm Dec 2011, Washington, DC. Sponsored License		• •			
14. ABSTRACT Chloroethenes, including vinyl chlorid analysis revealed that the vinyl chlorid embedded in a horizontally-acquired g similarity between vcrABC sequences horizontal acquisition driven by ~100 y most rdh genes relevant for bioremedithis mechanism. This talk will discuss molecular tools for monitoring and predehalogenases.	le reductase operon, genomic island that i from diverse Dehald years of industrial peation appear to be pothe implications of the second secon	vcrABC, of Dehantegrated at the sococooldes is quantellution with chlored redominantly acchese finding for description.	alococcoides a single-copy g ntitatively con orinated ether quired by De development	sp. strain VS is ene ssrA. The high nsistent with recent nes. Moreover, halococcoides via and use of	
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON	

Same as

Report (SAR)

c. THIS PAGE

unclassified

a. REPORT

unclassified

b. ABSTRACT

unclassified

25

GENOME-ENABLED MOLECULAR TOOLS FOR REDUCTIVE DEHALOGENATION

PROFESSOR ALFRED SPORMANN

Stanford University
Departments of Chemical Engineering, and of Civil & Environmental Engineering
318 Campus Drive
Stanford, CA 94305
(650) 723-3668
spormann@stanford.edu

Chloroethenes, including vinyl chloride, are widespread groundwater pollutants. A comparative genomic analysis revealed that the vinyl chloride reductase operon, vcrABC, of Dehalococcoides sp. strain VS is embedded in a horizontally-acquired genomic island that integrated at the single-copy gene ssrA. The high similarity between vcrABC sequences from diverse Dehalococcoides is quantitatively consistent with recent horizontal acquisition driven by ~100 years of industrial pollution with chlorinated ethenes. Moreover, most rdh genes relevant for bioremediation appear to be predominantly acquired by Dehalococcoides via this mechanism. This talk will discuss the implications of these finding for development and use of molecular tools for monitoring and predicting in situ remediation of chloroethenes via anaerobic reductive dehalogenases.








The 'old' paradigm

Bioremediation is mediated by a <u>single</u> important gene in a <u>single strain</u> that is <u>robust</u> to environmental perturbations



= Dehalococcoides sp.

= = reductive dehalogenase genes
















Dehalobacter sp.
Desulfitobacterium sp.
Dehalospirillum sp.
Geobacter sp.
Dehalococcoides sp.











Dehalobacter sp.
Desulfitobacterium sp.
Dehalospirillum sp.
Geobacter sp.
Dehalococcoides sp.


















16S rRNA gene-based phylogeny










Evolution of reductive dehalogenation traits in Dehalococcoides sp.



Reductive dehalogenase genes move as mobile elements in *Chloroflexi* populations









The 'old' paradigm

The new paradigm





= Dehalococcoides sp.


= reductive dehalogenase genes







Dehalobacter sp.
Desulfitobacterium sp.
Dehalospirillum sp.
Geobacter sp.
Dehalococcoides sp.










Dehalobacter sp.
Desulfitobacterium sp.
Dehalospirillum sp.
Geobacter sp.
Dehalococcoides sp.









How to assess gene dynamics and H₂ flux in heterogeneous environments?








How to assess gene dynamics and H₂ flux in heterogeneous environments?

Design of a tiled functional gene array



Hydrogenase gene array (H₂ase Chip)

~2300 diverse hydrogenase genes

Reductive dehalogenase gene array (Rdh Chip)


~280 diverse *rdh* genes







H₂ flux in complex environments










H₂ flux in complex environments



Hydrogenase chip can predict H₂ flux in complex communities








How do different (indirect) H₂ donor effect Dhc populations?













How do different (indirect) H₂ donor effect Dhc populations?







Dynamic changes in *Dehalococcoides* populations upon change in <u>H₂ source</u>








The new paradigm

Bioremediation is largely due to activity of mobile genes in heterogeneous populations that exhibit fluctuations upon environmental perturbations



= Dehalococcoides sp.

= = reductive dehalogenase genes








How do competing electron acceptors effect Dhc populations?









How do competing electron acceptors effect Dhc populations?







Poster 214




Dynamic change in *Dehalococcoides* population upon <u>alternate acceptor addition</u>







PM5L chemostat batch cultures (sulfide vs. sulfate addition experiments)




Sulfide addition is sufficient to explain Dehalococcoides population shift







Useful molecular tools reveal a shift in paradigm for reductive dehalogenation in complex environments



The new paradigm

Chloroethene bioremediation is largely due to dynamic, evolving, and heterogeneous populations of dehalogenating *Chloroflexi* that respond differentially to environmental perturbations.







Acknowledgement

Stanford group
Ian Marshall
Joey McMurdie

Koshlan Mayer-Blackwell

Sebastian Behrens Spormann group

Susan Holmes

Oregon State group
Lew Semprini
Mohammed Azizian

Frank Löffler, U Tenn/ORNL Elizabeth Edwards, U Toronto





